

1           Advanced prediction of the Intra-Americas Sargassum Season through Analysis of the  
2                           Sargassum Loop System Using Remote Sensing Technology

3  
4   Jeff Frazier

5   jtfrazier@email.tamu.edu

6   Co Author Dr. Tom Linton

7   Co Author Robert Webster

8   Abstract

9   Sargassum is a common type of seaweed observed in the Sargasso Sea, located in a portion of  
10 the western mid-Atlantic. Seasonally Sargassum inundates the beaches of Texas and the cost for  
11 its removal results in great strain on the coastal economies. Although this is an annual occurrence  
12 its cyclic migration patterns are relatively unknown. The research reported herein investigates  
13 the following null hypothesis, that Sargassum does not enter the Gulf of Mexico via the northern  
14 passages of the Caribbean and Yucatan Strait, where it amasses on the shores of the Gulf Coast  
15 or gets carried out the Florida Strait, in what is known as the Sargassum loop System. Once a  
16 seasonal migration patter is discerned, it is then hypothesized that certain aspects of the  
17 upcoming Sargassum season can be predicted in advance, using satellite imagery to monitor the  
18 corridors between the Sargasso Sea and the Gulf of Mexico.

19 The Sargassum season was previously thought to be erratic and unpredictable, however the  
20 theory of the Sargassum loop system sheds light on the seasonal migration patterns of the macro-  
21 algae. Through use of NASA's Landsat satellite imagery the presence and abundance of  
22 Sargassum has been analyzed. Based on several factors, such as ocean currents, wind patterns,  
23 time of the year, and size of seaweed mats, the arrival and intensity of the upcoming Sargassum

24 season can be approximated prior to its arrival in the Sargassum loop system. The Sargassum  
25 season starts months in advance in the Sargasso Sea when high pressure anomalies form. Their  
26 formation creates circulating northern wind currents that direct Sargassum southbound into the  
27 Caribbean latitudes. Once Sargassum has entered the Caribbean Passages, the Gulf Stream  
28 carries it westward, where depending on the direction and magnitude of ocean and wind currents,  
29 the Seaweed can take from two to five months to reach the Texas Coast. The Sargassum that  
30 does not reach land is flushed out through the Florida Straits and returns to the Sargasso Sea.  
31 This creates the Sargassum loop system. This, if monitored correctly, assists in forecasting the  
32 upcoming Sargassum season. Remote sensing, along with data from other ocean monitoring  
33 devices provides the necessary data for use in a Sargassum Early Advisory predictive model that  
34 allows for a more advanced warning of its arrival.

#### 35 Additional Key Words

36 Satellite Imagery, Sargasso Sea, Sargassum Early Advisory System, Azores High Pressure  
37 System

#### 38 Introduction

39 Sargassum is a free floating brown algae that has for thousands of years washed ashore on the  
40 coast of Texas during the summer months. Still, little is known about the path it takes to reach  
41 the coasts of the Western Gulf of Mexico. For centuries, Sargassum has been looked on as a  
42 plague because of its unsightliness and pungent odor produced by its decay. However, recent  
43 discoveries suggest that Sargassum mats deliver several benefits that are essential to the health of  
44 the oceans' and coastal ecosystems. These recent findings are revealing more positive aspects of  
45 Sargassum and are turning the tide on its historically negative image.

46 Until recently, there has never been a way to predict where Sargassum is going to land and in  
47 what quantities. This creates a bigger issue because there are not enough resources to sufficiently  
48 provide the entire coast with adequate Sargassum mitigation equipment during the peak of the  
49 Sargassum season.

50 In 2010, principle investigator Dr. Thomas Linton and Doctoral candidate Robert Webster  
51 initiated the Sargassum Early Advisory System (SEAS) program. It was a program, in which,  
52 through use of remote sensing technology, it would be possible to locate Sargassum mats  
53 (wrack) off the Texas Coast and predict, well in advance, when it would make landfall using  
54 remote sensing technology. By examining current satellite imagery from NASA's Landsat  
55 database, the SEAS program monitored for and analyzed Sargassum patterns in the Gulf of  
56 Mexico. Using oceanic and wind vector data, the program was able to create an approximate set  
57 and drift for the mats. These techniques allowed the SEAS program to effectively forecast the  
58 2011 Sargassum season and send out advisories predicting the landfall of Sargassum mats with a  
59 success rate of 84%. A successful advisory is defined as predicting a significant Sargassum  
60 wrack (enough to form a windrow) on the beach arriving within an eight day period. Before the  
61 SEAS program began, coastal communities had no means of determining where Sargassum  
62 would make landfall, nor in what volume. This caused the Sargassum relocation equipment to be  
63 spread thinly across the coast. This caused delays in response time. The advisories alerted coastal  
64 managers of the size and approximate location of an upcoming Sargassum episode, so that they  
65 could allocate and concentrate their Sargassum mitigation efforts more efficiently.

66 Satellite imagery limitations restricted the SEAS program's advisories to a maximum forecast  
67 period of sixteen days. Further, the satellite imagery did not detail the initiation and the intensity  
68 of the upcoming Sargassum season until it had arrived. In order to provide a more robust

69 advisory, several months in advance of the season, the SEAS program looked to expand its area  
70 of observation.

71 An atypical and unusually massive Sargassum event in the Southern Caribbean in 2011 initiated  
72 the expansion of the SEAS program, expanding its monitoring efforts to the Caribbean and  
73 Sargasso Sea. This paper theorizes that the observations in historical imagery of the newly  
74 incorporated region reveal a seasonal pattern of Sargassum migration. Annually, Sargassum is  
75 witnessed migrating south out of the Sargasso Sea and into the Caribbean, where it is swept  
76 northwest into the Gulf of Mexico by the Gulf Stream. Here it is theorized that Sargassum either  
77 washes ashore on the western coasts of the Gulf or is swept back out into the Atlantic, carried by  
78 the Gulf Stream current. This yearly voyage has been deemed the Sargassum loop current  
79 (Figure 1). The research reported in this paper attempts to document this Sargassum loop system  
80 so that it can be monitored and analyzed using remote sensing technology. This in turn, allows  
81 for an advanced forecast of the upcoming Intra-American Sargassum season.

82 Since Sargassum rests atop of the water column, the two main forces that affect it are ocean  
83 currents and wind currents. The effects of the ocean currents appear more apparent because they  
84 are well mapped and tend to be static in location, direction, and intensity. The Sargasso Sea is  
85 formed by the circulation of the North Atlantic Gyre. Sargassum accumulates in the center of the  
86 Gyre as a result of this phenomenon, until other, more prevailing forces affect its location. Wind  
87 currents have a surprisingly significant impact on Sargassum. With no freeboard, one would  
88 expect Sargassum to have little response to atmospheric conditions, however, this study suggests  
89 that the initiation of Sargassum migration is most closely correlated with the Azores High  
90 Pressure System. This high pressure weather phenomenon generates southbound winds over the  
91 Sargasso Sea, causing the Sargassum to migrate into lower latitudes. The mechanism leading to

92 this magnitude of effect by wind intensity and direction on Sargassum, is still being studied,  
93 however the distinct correlation of wind patterns and seaweed migration can be inferred.

94 The main force that drives wind is known as an atmospheric phenomenon called the North  
95 Atlantic Oscillation occurs over the Northern Atlantic Ocean. In this phenomenon, a series of  
96 pressure systems, alternating between high pressure and low pressure, move across the northern  
97 hemisphere of the Atlantic Ocean (Figure 2).

98 When a high pressure system has moved into the region of interest, it is said to be in a positive  
99 mode. The high pressure systems are characterized as having a mean sea level pressure of at least  
100 1010 millibars, but commonly range between 1020 to 1030 millibars, and is anticyclonic,  
101 meaning it rotates clockwise. One particular high pressure system situated in this region is  
102 commonly referred to as Azores High Pressure System, and oscillates from the Eastern to the  
103 Western Atlantic, frequently around latitudes of 30 degrees.

104 The Azores High Pressure System is theorized to be the initiating force behind Sargassum  
105 migration from the Sargasso Sea to the Gulf of Mexico. During the winter and spring months, the  
106 Azores Pressure System shifts over to the Eastern Atlantic where it resides over the Sargasso Sea  
107 (Figure 2). Anti-cyclonic conditions create swift northwesterly winds over the Sargasso Sea.

108 These winds are believed to be a strong enough driving mechanism to push the floating algae out  
109 of the North Atlantic Gyre and down south toward the Caribbean, initiating the Sargassum Loop  
110 System.

111 During the summer season, a Sargassum accumulation off the western coast of the Gulf of  
112 Mexico creates a second Sargassum reservoir, from which the seaweed washes ashore in massive  
113 quantities. Such occurrences have been documented, in newspapers, on the coast of Texas dating  
114 back to the late eighteen hundreds. While the intensity of the summer Sargassum episodes have

115 varied greatly from year to year, it has continued to occur, without explanation. Until now, little  
116 was known about the driving mechanism and origin of this seasonal Sargassum cycle. The  
117 research findings set forth in this paper describe the studies undertaken to explain the underlying  
118 mechanism by which the annual season of Sargassum occurs, as well as the area of origin of the  
119 wracks seen on the Texas Gulf Coast.

120 In 2009, Dr. Thomas Linton and Doctoral candidate Robert Webster formed the Sargassum Early  
121 Advisory System (or SEAS). The SEAS program set out to predict upcoming Sargassum events  
122 on the Texas Coast, utilizing remote sensing technology. Using satellite imagery, the coastal  
123 waters of Texas were monitored and analyzed for presence of Sargassum (Webster 2014).  
124 By 2012, the SEAS program had refined its process of forecasting seaweed episodes to provide  
125 up to 97% accuracy. However, the most advanced warning of an impending episode an advisory  
126 could give at the time was sixteen days (Webster 2014). The mechanism that initiates the  
127 Sargassum season on the Texas Coast had yet to be discovered. Once this was determined, it was  
128 theorized that entire Sargassum seasons could be forecasted, giving unprecedented forewarning  
129 to the coastal communities in Texas.

130 So where does Sargassum originate from and where does it congregate during winter months?  
131 One source suggests that a reservoir of Sargassum forms in the Bay of Campeche annually  
132 during the winter months, then drifts north and amasses in spring and summer months,  
133 depositing on the shores of the Gulf of Mexico (Gower and King 2012). A review of historic  
134 satellite imagery showed no observable amounts of Sargassum in that area during the winter  
135 months. The concept of Sargassum originating in the Sargasso Sea, along the Mid Atlantic Bight,  
136 seemed evident; however the path from the East coast of the United States to the Western Gulf of  
137 Mexico appeared much less certain. The path from the Atlantic through the Florida Straits to the

138 Gulf was the most direct. However a close examination of the currents between Florida and Cuba  
139 shows that the swift Gulf Stream is heading easterly and directly opposing the route which  
140 Sargassum must take to get to the Western Gulf of Mexico.

141 In 2011, an atypical event occurred in the Southern Caribbean islands that initiated the idea that  
142 Sargassum may drift as far south as the Caribbean from the Sargasso Sea. It was theorized that if  
143 Sargassum could seasonally drift to tropical latitudes, then it could utilize westerly Caribbean  
144 currents to enter the Gulf through the Yucatan Passage. But if this event in 2011 was so  
145 uncommon, how could a seasonal occurrence of Sargassum in the Caribbean go unnoticed for so  
146 long? An in-depth examination of the 2011 Sargassum episodes in the Southern Caribbean  
147 revealed the connection between the Sargasso Sea and the seasonal Sargassum in the Gulf of  
148 Mexico.

#### 149 Analysis

150 The 2011 Southern Caribbean Sargassum event generated the theory that Sargassum has the  
151 ability and resiliency to migrate from the Sargasso Sea into tropical waters. However, since  
152 Sargassum sightings have been sparsely recorded in the historical archives of the Southern  
153 Caribbean, one can deduce that an irregular occurrence sparked this 2011 phenomenon. This  
154 means that Sargassum could potentially migrate south annually, however, under normal  
155 conditions, it won't propagate as it did in 2011. So what happens to the Sargassum once it enters  
156 the Caribbean Sea?

157 Hypothesis 1) Annually, atmospheric wind currents drive Sargassum south out of the Sargasso  
158 Sea into the Northern Caribbean, where the Gulf Stream carries it into the neritic Gulf of  
159 Mexico. Once in the Gulf, it amasses and is either carried west by prevailing wind currents or

160 carried out the Florida Strait back to the Sargasso Sea via the Gulf Stream, in what is known as  
161 the Sargassum loop system.

162 The Azores high pressure system produces concentrated northern winds across the Sargasso Sea,  
163 which pushes a portion of the sea of macro-algae out of the North Atlantic Gyre and into the  
164 Caribbean. Once in the Caribbean, the anti-cyclonic wind currents assist the Gulf Stream in  
165 carrying the Sargassum through the Yucatan Strait and into the Gulf of Mexico. Sargassum is  
166 then carried through the loop current, where wind or fluctuations in the loop current can cause  
167 the algae to break off and drift westward, where it will land on the Texas Coast. The remaining  
168 Sargassum that does not deviate from the loop current is carried back out into the Mid Atlantic  
169 bight, where it returns from its Intra-Americas journey.

170 In order to investigate the theory of the Sargassum loop system, The SEAS program expanded  
171 their area of observation to the northern Caribbean and southern Sargasso Sea regions. Using  
172 historic satellite imagery, the Sargassum loop system can be observed as it occurs annually  
173 throughout the Intra-Americas. An examination of this archived imagery reveals several patterns  
174 that occur seasonally and could be an earlier indication of the Sargassum migration, allowing for  
175 a more advanced prediction of the impending Sargassum season.

176 Hypothesis 2) The Impending Sargassum season for the Intra-Americas can be predicted in  
177 advance by comparing migration patterns of Sargassum in current satellite imagery to the  
178 migration patterns of Sargassum in historic satellite imagery.

179 Satellite imagery with sufficient resolution to see Sargassum is archived as far back as 2000.

180 During the period of January of 2000 to May of 2013, the Landsat satellites created an archive of  
181 347 images for each passage, not accounting for any imagery that is unusable, such as complete  
182 cloud cover, or data missing from the satellite imagery database. In order to obtain an



183 appropriate sample of images of the passages, 70 images (or 20 percent of the archived database)  
184 of each passage were analyzed for the presence of Sargassum (figures 3a and 3b for example).  
185 Sargassum manifests in two forms. If the pelagic mat is large enough, it will produce a green  
186 glow in the satellite image, this is due to the vegetative growth filter added to the landsat images.  
187 The second manifestation of Sargassum is what is referred to as a 'slick' meaning that  
188 Sargassum itself cannot be seen in the image, however its presence at the ocean surface disrupts  
189 the surface tension and wave attenuation, thus creating a darker slick around the otherwise  
190 invisible mat. An examination of archived satellite imagery of the newly expanded area of  
191 observation revealed that not only has Sargassum been historically present, but also that it was  
192 seen in patterns that appeared to repeat annually.

193 Several patterns involving the southern migration of Sargassum emerged from the historical  
194 analysis of Caribbean Satellite imagery. The first pattern involves the initiation of Southern  
195 migration of Sargassum which starts as early as December, but typically occurs between  
196 February and April (See Table 1). Sargassum can be seen breaking away from the typical  
197 boundaries of the Sargasso Sea set by the motion of the North Atlantic Gyre. It is theorized that  
198 seasonal northerly winds produce adequate locomotion for the Sargassum to break away from its  
199 origin, pushing it south, toward the Caribbean. Although gyrating currents formed from the  
200 Coriolis Effect are strong enough to concentrate the Sargassum into the center of the gyre, these  
201 forces are relatively weak compared to locomotive forces produced by the wind and the Gulf  
202 Stream. Even though Sargassum has little to no freeboard, wind currents still affect Sargassum  
203 drift significantly. The estimated time it takes to get from the Sargasso Sea to the Gulf Coast is  
204 roughly two to five months, so the long range prediction indicates the initiation, intensity, and  
205 climax of the upcoming Sargassum season. This is shown in Table 1, which lists the number of

206 sightings of Sargassum seen in each month. A spike in observations occurs during the winter to  
207 spring months (March through May). This precedes the Sargassum episodes seen on the Texas  
208 Coast by two to five months. For accurate and precise forecasting, the prediction of landfall is  
209 not made until after the Sargassum has entered the Gulf of Mexico.

210 Once it has broken free from the North Atlantic gyre, a second pattern begins to manifest. As the  
211 free-floating algae reaches the Northern Caribbean Islands, it either washes ashore on the islands  
212 of Cuba, Haiti, Dominican Republic, and/or Puerto Rico, or it gets funneled through the passes  
213 between the islands and into the Caribbean Sea (See Figures 3 and 4). Because of the passage are  
214 narrow, monitoring the Windward, Mona, and Anegada Passages are emphasized.

215 Once Sargassum has reached the neritic waters of the Caribbean Sea, the chemistry of the water  
216 is more favorable for Sargassum growth. The shallow, coastal waters have relatively high  
217 concentrations of nutrients such as carbon and phosphorous, compared to the nutrient poor  
218 waters of the Atlantic. Once in the Caribbean, the Sargassum mats start to increase in size.

219 The third and final pattern that emerged from the examination of the historical Caribbean  
220 satellite imagery is observed once the Sargassum has reached the Caribbean. Sargassum is seen  
221 migrating westerly once in the Caribbean Sea (Figure 5). An analysis of the Gulf Stream shows a  
222 relatively concentrated current from east to west in the Caribbean Sea, which is theorized to  
223 assist Sargassum in its annual migration (Figure 6). It is observed exiting the Caribbean through  
224 the Yucatan Passage where it arrives in the Gulf of Mexico. This seasonal Sargassum migration  
225 theory connects the Sargasso Sea, or the origin, to the Gulf, where it can be seen deposited along  
226 the coastline of Texas.

227 This annual voyage that Sargassum makes has been called the Sargassum loop system. During  
228 the winter months, atmospheric conditions create northerly winds that initiate Sargassum's

229 yearly migration south, through the Northern Caribbean passage (Table 1). Once it reaches the  
230 Caribbean, it is caught in the Gulf Stream, where it is taken by concentrated surface currents into  
231 the Gulf of Mexico.

232 The discovery of this Sargassum loop system now allows for the monitoring, analysis, and long  
233 range forecasting of the Texas' annual Sargassum season via remote sensing. Since seaweed that  
234 eventually lands on the Texas coast has drifted south through the Caribbean passages several  
235 months prior, one can use this foresight to advise the coastal communities of the forecast for the  
236 initiation of the upcoming Sargassum season, as well as the intensity of the Sargassum episodes  
237 expected to occur that season, and the commencement of landfall events.

238 The start of the Sargassum season occurs several months after atmospheric conditions allow  
239 Sargassum to migrate south out of the Sargasso Sea and into the Caribbean. Since there is still a  
240 distance of over 3,000 miles to be traveled, several factors can affect the Sargassum before it  
241 makes landfall on the Texas coast. Variables, such as oceanic and atmospheric conditions are  
242 dynamic, growth rates are affected by nutrient and temperature conditions during the migration,  
243 which can affect the quantity of Sargassum in the Sargassum loop system, as it amasses. Not all  
244 Sargassum that journeys into the Gulf will make landfall on the coast, but instead be discharged  
245 through the Florida Straits. Because of these reasons, at this point in the loop system, exact  
246 landfall dates cannot be predicted.

247 Relative correlations between the amount of Sargassum that can be seen in the passages and the  
248 amount that makes landfall can be made. It is observed that Sargassum is more commonly  
249 sighted in the passages during seasons of heavier Sargassum inundations, such as the 2008 and  
250 2011 Sargassum seasons, of the coastal communities of the Gulf (Table 2). In contrast, during  
251 lighter years of Sargassum episodes, such as the 2006 and 2010 Sargassum seasons, Sargassum

252 sightings in the Caribbean passages occur less commonly (Table 2). The frequency of Sargassum  
253 sightings in the passages can be directly correlated with the abundance of Sargassum in the Gulf  
254 and therefore, can forecast the intensity of the upcoming season, however one cannot extrapolate  
255 an absolute volume.

256 The end of the Sargassum season occurs when atmospheric conditions that encourage Sargassum  
257 become reduced in frequency and intensity, reducing the locomotive energy in the wind, the  
258 driving force of the Sargassum migration. This termination of the seaweed season is observed in  
259 both the degradation of the northerly winds over the Sargasso Sea and in the absence of  
260 Sargassum in the Caribbean passages.

261 There are a few caveats in the advanced prediction of the upcoming Sargassum season. First, the  
262 model used to predict the set and drift of flotsam and jetsam (floating debris) in the Gulf of  
263 Mexico is still in the experimental phase. Several entities have attempted to create drift models,  
264 such as NASA's debris prediction model, Coast Guard's Oil prediction model, and NOAA's  
265 trash and flotsam predictive model, however none have proven accurate or effective for  
266 Sargassum prediction or for their original intended purposes. This forces us to use a more  
267 simplistic and experimental model of vector addition. Once Sargassum is located, the regional  
268 ocean and wind currents are obtained and overlaid over the mat of Sargassum. It is known that  
269 wind has a stronger effect on Sargassum than ocean currents, despite Sargassum having a  
270 negligible amount of freeboard, allowing it to be more affected by air. Because of this, the wind  
271 current is given more weight when modeling the drift of the seaweed mats. The exact effect that  
272 wind and water currents have on Sargassum is still unknown. However. It is noted that flotsam  
273 and jetsam travel at a rate of 3 percent of the currents acting upon it (Gyory 2013). i.e. If ocean  
274 and atmospheric currents are acting on Sargassum at a rate of 15 knots, the Sargassum mat is

275 only drifting at a rate of 0.45 knots. This prediction model is simplistic and is still in the early  
276 experimental stages of development.

277 Another constraint on long range forecasting is provided by the dynamic variables. Since  
278 Sargassum is being forecasted months in advance, there are several factors such as ocean and  
279 atmospheric conditions that change daily, and cannot be accurately incorporated into the model.  
280 Because of this, a more simplistic average is being taken of the wind and ocean currents. Another  
281 variable that affects the drift of Sargassum is the bathymetry, which affects wave patterns and  
282 tidal cycles. Since Sargassum must travel a minimum of 1,800 miles, accounting for the  
283 bathymetry and the added variables that it entails, it is impractical to incorporate them into the  
284 model. These discrepancies in the modeling for Sargassum set limitations on the accuracy of  
285 Sargassum predictions.

286 Expanding the SEAS program's area of observation into the Caribbean has increased the  
287 forewarning time from sixteen days to up to two to five months in advance. Knowledge of the  
288 initiation and intensity of the upcoming summer seaweed episodes allows coastal communities to  
289 allocate the necessary resources and better prepare for the impending Sargassum season.

## 290 Conclusion

291 The increased foresight brought on the by expanding the SEAS's area of observation into the  
292 Caribbean, allows for the forecasting of Sargassum in the Gulf of Mexico months in advance. It  
293 also helps explain the secret to the sustainability of the Sargasso Sea, an oasis of vegetation  
294 located in a nutrient-barren region. The 2011 massive Sargassum event in the Southern  
295 Caribbean was then thought to be generated by unusually massive flooding. Although it may  
296 have seemed like a dead end, this investigation of the unusually massive Sargassum episode led  
297 to the theory of Sargassum's annual migration from the Sargasso Sea to the Gulf and the creation

298 of the Long range forecasting of Sargassum in the Caribbean Sea. Previously, Sargassum could  
299 not be forecasted until it had reached the coastal waters of the Gulf of Mexico. This expansion of  
300 the SEAS program allows for the upcoming Sargassum season on the Texas Coast to be  
301 forecasted months in advance. The frequency of appearance and volume of Sargassum in the  
302 Caribbean passages during the winter and spring months are indications of the initiation,  
303 intensity and eventually the commencement of Sargassum season in the Caribbean and Gulf of  
304 Mexico.

305

#### 306 References

307 Gower, Jim. & King, Stephanie. Satellites Suggest a New Source of Caribbean Sargassum.  
308 British Columbia, Canada. Fisheries and Oceans Canada, Institute of Ocean Sciences. 2012.  
309 Print.

310

311 Gyory, J., A.J. Mariano, and E. H. Ryan. "The loop current." Ocean Surface Currents. 2013.  
312 Web. 02 Apr. 2014. <http://oceancurrents.rsmas.miami.edu/atlantic/loop-current.html>

313

314 HYCOM. "Ocean Prediction: Intra-Americas Snapshot Archive." Hybrid Coordinate Ocean  
315 Model. Center for Ocean-Atmospheric Prediction Studies, 4 Mar. 2014. Web. 11 Apr. 2014.  
316 <http://hycom.org/>.

317

318 Masters, Jeff. "Dr. Jeff Masters' WunderBlog." March Like a Lion: Record Cold in Midwest,  
319 Heavy Snow in Mid-Atlantic. Weather Underground, 01 Mar. 2014. Web. 03 Mar. 2014.  
320 [http://www.wunderground.com/data/640x480/huratl\\_slp.gif#Mean%20Sea%20Level](http://www.wunderground.com/data/640x480/huratl_slp.gif#Mean%20Sea%20Level)

321

322 Texas Automated Buoy System. Vector addition trajectory tool. Texas General Land office,

323 Geochemical & Environmental Research Group. May, 2014. Web. 02 June, 2014.

324 <http://tabs.gerg.tamu.edu/>

325

326 United States Geological Survey. "USGS Global Visualization Viewer." USGS Global

327 Visualization Viewer. United States Geological Survey, 02 Mar. 2014. Web. 03 Mar. 2014.

328 <http://glovis.usgs.gov/>

329

330 United States Geological Survey. "EarthExplorer." EarthExplorer. United States Geological

331 Survey, 02 Mar. 2014. Web. 03 Mar. 2014. <http://earthexplorer.usgs.gov/>


332

333 Webster, Robert. "Sargassum Early Advisory System." Sargassum Early Advisory System

334 (SEAS). Texas A&M at Galveston, 14 Feb. 2014. Web. 03 Mar. 2014. [www.tamug.edu/SEAS](http://www.tamug.edu/SEAS)

335

336 Figures and Tables




337

338  
339  
340  
341  
342  
343  
344

**Figure 1 Sargassum loop system**-This graphic represents the *Sargassum* loop system that initiates in the Sargasso Sea, as developed by the SEAS program. Atmospheric conditions create wind patterns that push *Sargassum* south, into the Caribbean where it is pushed west, by the oceanic and atmospheric currents carrying it into the Gulf of Mexico. There it either washes ashore on the Gulf Coast or gets swept out the Florida Strait via the Gulf Stream (Webster, 2014)






**Mean Sea Level Pressure and 1000 mb Winds  
6 hour GFS Forecast Valid 18 GMT Sun Mar 02**

345  
346  
347  
348  
349

**Figure 2 Azores High Pressure System Graphic–The Azores High Pressure system creates anticyclonic winds over the Sargasso Sea and becomes the driving mechanism behind the migration of *Sargassum* (Masters, 2014)**



350

351 **Figure 3a Mona Passage Satellite Image**—This graphic illustrates a satellite image  
352 **of *Sargassum* migrating through the Mona Passage into the Caribbean Sea.**  
353 ***Sargassum* and its slicks are denoted by the yellow oval (United States**  
354 **Geological Survey, 2014).**  
355



356

357 **Figure 3b Mona Passage Satellite Image Magnified**—A magnified view of these  
 358 images reveal a linear green glow or if the mat is too small to be seen in the  
 359 image, a darker outline in linear patterns. This is because the size of the mat is  
 360 too small to be seen in the image, only the disturbance of the wave attenuation  
 361 around it can be seen (United States Geological Survey, 2014).


362



363

364 **Figure 4 Caribbean Passages** – This figure shows a map of the Caribbean. The  
 365 Northern Caribbean Passages, referred to as the Windward, Mona, and Aneгада  
 366 Passages, are where *Sargassum* is most frequently observed (Webster, 2014).


367



368  
 369  
 370  
 371  
 372  
 373  
 374

**Figure 5 Yucatan Strait Satellite Image – A satellite image of the Yucatan Strait on March 22, 2014 illustrating *Sargassum* moving westward via the Gulf Stream. *Sargassum* manifests itself in the form of a slick. Since the size of the mat is too small to be seen in the image, only the disturbance of the wave attenuation around it can be seen (United States Geological Survey, 2014).**

**Speed/currents layer 1 Oct 31, 2013 00Z 91.0**




375

376  
 377


**Figure 6 Gulf Stream Current Graphic-This graphic represents the Gulf Stream that generates a western current in the Caribbean. The current is then pushed**

378 north into the Gulf of Mexico where it 'loops' and exits out of the Gulf via the  
379 Florida Strait (HYCOM, 2014).  
380




381

382 **Figure 12 Florida Strait Satellite Image**—a Satellite image of the Florida Strait On  
383 **April 24, 2014** illustrates *Sargassum* exiting the Gulf of Mexico via the Florida  
384 **Strait** and rejoining the Sargassum Sea (United States Geological Survey, 2014).  
385



386  
 387 **Table 1 *Sargassum* Sightings in the Caribbean Passages**—This graphic illustrates  
 388 the number of times *Sargassum* has been spotted in the passages in a single  
 389 month. Historic satellite imagery was obtained and analyzed to review when  
 390 *Sargassum* begins its journey through the *Sargassum* loop system.  
 391



392  
 393 **Table 2 Intensity of *Sargassum* Sightings per Year**—This graphic depicts the  
 394 annual difference between *Sargassum* sightings in the Caribbean Passages. Note  
 395 that during the 2008 and 2011 seasons, more *Sargassum* was seen the Caribbean  
 396 Passages, which correlates to the relatively higher rates of *Sargassum* events on  
 397 the Texas Coast. The 2007 and 2010 years yielded less *Sargassum* in the  
 398 passages, which correlates to the lower volumes of *Sargassum* seen in the Gulf  
 399 of Mexico. This denotes that the intensity of the upcoming *Sargassum* season is  
 400 reflective in the Caribbean Passages during precluding months.  
 401